



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

It's well and truly time to stop stating that AMPK regulates glucose uptake and fat oxidation during exercise

This is the Accepted version of the following publication

McConell, Glenn (2020) It's well and truly time to stop stating that AMPK regulates glucose uptake and fat oxidation during exercise. *American Journal of Physiology - Endocrinology and Metabolism*, 318 (4). E564-E567. ISSN 0193-1849

The publisher's official version can be found at
<https://journals.physiology.org/doi/abs/10.1152/ajpendo.00511.2019>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/41616/>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

**It's well and truly time to stop stating that AMPK regulates
glucose uptake and fat oxidation *during* exercise**

Glenn K. McConell

Institute for Health and Sport, Victoria University, Melbourne, Victoria, 8001, Australia

Running title: Does AMPK really regulate exercise metabolism?

Corresponding author:

Glenn K McConell

Institute for Health and Sport (IHes), Victoria University

PO Box 14428, Melbourne

VIC 8001, Australia

Email: glenn.mcconell@vu.edu.au

Phone: +613 9919 9472

Fax: +613 9919 9480

AMPK and exercise

In the late 1990s it was shown *in vivo* that skeletal muscle AMP-activated protein kinase (AMPK) activity is increased during exercise whilst pharmacological activation of AMPK could also increase skeletal muscle glucose uptake and free fatty acid (FFA) oxidation *in vivo* (22, 37). Taken together, these findings provided the impetus for the multitude of subsequent studies that have been undertaken to elucidate the role of AMPK in skeletal muscle metabolism. AMPK is a $\alpha\beta\gamma$ heterotrimer comprised of two α , two β , and three γ subunits (8). The α -subunits contain the catalytic domain responsible for activation which, during contraction, occurs via an increase in cellular stress due to increases in AMP_{free} and ADP_{free} levels. The increase in AMP_{free} allosterically activates AMPK, makes AMPK more susceptible to phosphorylation by upstream AMPK kinase(s), and the increase in AMP_{free} and ADP_{free} antagonizes the effect of protein phosphatases on AMPK, reducing the likelihood of dephosphorylation of AMPK (8, 40). AMPK is activated during moderate or harder exercise in humans (3, 6, 39) and there is evidence that AMPK phosphorylates and inactivates acetyl-CoA carboxylase (ACC)- β which is then thought to increase fat oxidation (37). However, AMPK cannot simultaneously act as a primary regulator of both glucose uptake and FFA oxidation during exercise because there are situations (e.g. increases in exercise intensity, pre- vs. post-training exercise, high vs. low carbohydrate diets) whereby increases in glucose uptake are accompanied by simultaneous decreases in fat oxidation, and vice-versa. Indeed, below I will discuss evidence that AMPK actually does not regulate either glucose uptake or fat oxidation during exercise.

AMPK is not required for skeletal muscle glucose uptake during exercise

Support for a role of AMPK in the regulation of skeletal muscle glucose uptake were initially derived from pharmacological studies whereby the cell permeable agent 5-aminoimidazole-4-carboxamide-1- β -O-ribofuranoside (AICAR), an AMP mimetic, was shown to increase AMPK activity, Glut-4 translocation and glucose uptake (16, 22). However, it is important to note that whilst AICAR and contraction both activate AMPK and increase glucose uptake, this appears to be mediated via different pathways. While skeletal muscle glucose uptake in response to hypoxia/AICAR appears to be highly dependent on AMPK (7, 13, 25), as explained below there is evidence that this is not the case for contraction-stimulated glucose uptake. Thus, attempting to compare between AICAR, hypoxia, and contraction studies in relation to AMPK and glucose uptake can be problematic.

Although skeletal muscle AMPK α 2 activity activation and glucose uptake go hand in hand with increases in exercise intensity in humans (2) and rats (27), one cannot simply assume that this indicates a cause and effect relationship. Indeed, most human and rodent studies have found clear dissociations between skeletal muscle AMPK activation and glucose uptake. Already in 2002 Wojtaszewski et al. (38) found increases in leg glucose uptake during the first hour of cycling at $\sim 45\%$ $\text{VO}_{2\text{peak}}$, yet muscle AMPK α 2 activity and AMPK α Thr172 phosphorylation only significantly increased at the end of exercise (~ 3.5 hours). In addition, glucose uptake at the end of exercise was actually lower than earlier in exercise (38). Also, whole-body glucose uptake and skeletal muscle AMPK

activity increase 8-10 fold during prolonged cycling exercise at ~66% VO_2 peak in untrained humans, but after just 10 days of exercise training there is no measurable increase in AMPK activity at all during the same absolute intensity of exercise in these individuals (19). The exercise training did attenuate the increase in glucose uptake during prolonged exercise, but it was still substantial, so if AMPK was playing a role one would expect some increase in skeletal muscle AMPK activity but none occurred (19). Similarly, there is no activation of skeletal muscle AMPK during 120 min of exercise at ~65% VO_2 peak in well trained endurance athletes (McConell et al. in review), despite substantial glucose uptake at this intensity in well trained individuals (34). It should be noted though that there may have been activation of some subunits of AMPK that were missed when only total AMPK activity is measured. Indeed, although increases in AMPK activity that occur before exercise training are totally nonexistent during exercise at the same relative intensity (~65% VO_2 peak) after 12 weeks of exercise training, there is still 20% of the pre-training increase in the $\text{AMPK}\alpha 2\beta 2\gamma 3$ trimer of AMPK during such exercise (24).

Whole-body deletion of AMPK $\alpha 1$ or $\alpha 2$ fails to inhibit contraction-stimulated glucose uptake *in vitro* (13), whilst muscle glucose uptake during *ex vivo* contraction is either normal or only partially reduced in mice expressing a dominant-negative AMPK $\alpha 2$ isoform (7, 23, 25, 35)). When interpreting these findings one should consider the potential confounding effects of knocking-out or suppressing AMPK or LKB1, such as alterations and/or compensation of enzymes upstream and/or downstream. For example, knocking-out AMPK $\alpha 2$ results in compensatory increases in AMPK $\alpha 1$ expression in

mouse skeletal muscle (15), whilst skeletal muscle of the AMPK $\alpha 2$ dominant-negative mouse has attenuated force generation *ex vivo* (7, 25). Importantly, muscle-specific AMPK $\alpha 1\alpha 2$ double knockout (mdKO) mice, which have only 3-4% of normal AMPK $\alpha 2$ activity and no activation of AMPK during exercise, have normal increases in skeletal muscle glucose uptake during exercise at the same relative intensity (14).

Some findings over the years have provided small indications that AMPK may play a role in glucose uptake with contraction in rodents but it is important to consider that these situations are often non physiological. For example, it was shown that glucose uptake was lower during *ex vivo* twitch contractions in AMPK $\alpha 1$ KO mouse muscle than wild type mouse muscle and it was suggested that this may indicate that AMPK $\alpha 1$ plays a role in glucose uptake during short-duration low intensity exercise (11). But AMPK $\alpha 1$ is usually not activated during short-duration low-moderate intensity exercise (40-70% VO_2 max) (6, 11, 26, 38, 39).

A recent transgenic mouse study by Kjobsted et al. (14) convincingly showed that AMPK is not required for glucose uptake during contraction/exercise but rather is important for metabolism after exercise. Using muscle specific AMPK $\alpha 1\alpha 2$ double KO and inducible muscle specific AMPK $\alpha 1\alpha 2$ double KO mice, that have extremely low AMPK activity, they showed that glucose uptake was normal during *ex vivo* and *in situ* contraction. They also mentioned in the discussion (“unpublished observations”) that glucose uptake is normal in these mice during exercise (14). The authors said “We cannot ultimately rule out that AMPK may be necessary for regulating muscle glucose uptake during longer

periods of exercise (>30 min) and contraction (>10 min) where the myocellular stress may be further elevated” (14). However, as mentioned above, dissociations between glucose uptake and AMPK activation have been observed in humans during exercise lasting up to 3.5 hours (19, 38).

An important point made by the authors of the recent rodent study (14) was that of the 10 studies conducted previously that found impaired glucose uptake with contractions in AMPK-deficient mouse muscle, 8 of these actually examined the glucose uptake after, rather than during the contractions. As we pointed out previously in regards to a similar situation in studies examining the role of nitric oxide in glucose uptake with contraction (18), such study designs more investigate the effect of the intervention on glucose uptake after, than during, exercise.

Therefore, it is clear from both mouse and human studies that AMPK is not required for glucose uptake during exercise. It is not possible however to discount it altogether however because there is likely redundancy in signalling to skeletal muscle glucose uptake during exercise.

AMPK is not required for skeletal muscle fat oxidation during exercise

In skeletal muscle, malonyl-CoA is believed to be a major factor involved in the regulation of FFA oxidation due to its inhibitory effects on carnitine palmitoyltransferase (CPT) 1, the main enzyme controlling fatty acid flux into the mitochondria for oxidation. Malonyl-CoA is synthesized by ACC- β , and skeletal muscle contraction and AMPK activation phosphorylates ACC- β at a serine residue (212 in rodents, 221 in humans) in skeletal muscle (2, 8, 31), and *in vitro* this phenomenon inhibits ACC- β (37). This has led

to the assumption that the activation of AMPK during exercise increases FFA oxidation due to AMPK phosphorylating and inhibiting ACC- β and thus reducing malonyl-CoA and removing the inhibition of CPT1. However, as with glucose uptake, many dissociations have been observed between AMPK activation, ACC- β phosphorylation, ACC- β activity and FFA oxidation during contraction and exercise.

Low intensity exercise performed for ~3.5 hours in humans progressively increases skeletal muscle net FFA uptake and fat oxidation, despite the fact that ACC- β phosphorylation is only increased at 1hr, and AMPK activity and is only elevated at the end of exercise (38). Similarly, low intensity contraction increases FFA oxidation in rat skeletal muscle despite no alterations in AMPK or ACC- β activity (32). It has also been shown in humans that increasing exercise from moderate- to high-intensity results in a suppression of fat oxidation, but AMPK activity and ACC- β phosphorylation both increase (2) and ACC activity decreases (4). Furthermore, 10 days of exercise training in previously untrained individuals augments fat oxidation during prolonged exercise, yet skeletal muscle AMPK activity was not increased at all and ACC- β phosphorylation only increased slightly unlike larger increases during exercise before training (19).

AMPK α 2 DN mice, despite having very low AMPK activity, have normal increases in FAT/CD36 translocation to the plasma membrane (12), palmitate uptake (12) and fat oxidation (5) during contraction/exercise. Surprisingly, they also have normal increases in ACC β phosphorylation (5) and reductions in malonyl CoA (5) during contraction. It has also been shown in perfused rat hindlimbs that fat uptake increases within one minute

of in situ contractions but AMPK is not activated until later (12). A study that did find a reduction in fat oxidation during ex vivo contraction of muscle specific AMPK $\alpha 1\alpha 2$ KO mice also found that these mice had reduced CD36 and FABPpm, important for fatty acid uptake and oxidation, which likely accounted for these findings (15). Finally, there is evidence from skeletal muscle ACC β Ser 212 knock in mice that ACC β is important for AICAR-activated fatty acid oxidation at rest (30), but ACC β is not required for fatty acid oxidation during ex vivo contraction or during treadmill exercise (29). Therefore, transgenic models support the human findings that there are dissociations between AMPK and fat oxidation during contractions and exercise.

Conclusions

In conclusion, although activation of AMPK during exercise appears to be critical for adaptations after exercise (10, 14, 17, 21, 28), AMPK activation is not necessary for increases in glucose uptake or fat oxidation during exercise. In 2005 George Brooks wrote “Now, with the results of McConnell *et al.* (19). it is time to reassess the relevance of the AMPK signalling pathway for the regulation of metabolism in working human muscle” (1). Despite this, 14 years later most papers related to AMPK still state in the introduction that AMPK regulates glucose uptake and fat oxidation during exercise. Although redundancy likely exists, it really is time to move on and look more closely at other possible regulators of glucose uptake and fat oxidation during exercise such as nitric oxide (20, 33), Rac1 (33, 36) and ROS production (9, 33).

Acknowledgments

The author would like to thank the National Health and Medical Council (NHMRC) of Australia and Diabetes Australia for funding. In addition, GKM was a Danish Diabetes Academy Visiting Professor during the writing of this manuscript (supported by a research grant from the Danish Diabetes Academy, which is funded by the Novo Nordisk Foundation, grant number NNF17SA0031406).

REFERENCES

1. Brooks GA. Governor recalled! Now what? *J Physiol* 568: 355, 2005.
2. Chen Z-P, Stephens TJ, Murthy S, Canny BJ, Hargreaves M, Witters LA, Kemp BE, McConell GK. Effect of exercise intensity on skeletal muscle AMPK signaling in humans. *Diabetes* 52: 2205-2212, 2003.
3. Chen ZP, McConell GK, Michell BJ, Snow RJ, Canny BJ, Kemp BE. AMPK signaling in contracting human skeletal muscle: acetyl-CoA carboxylase and NO synthase phosphorylation. *Am J Physiol Endocrinol Metab* 279: E1202-1206, 2000.
4. Dean D, Dugaard JR, Young ME, Saha A, Vavvas D, Asp S, Kiens B, Kim KH, Witters L, Richter EA, Ruderman N. Exercise diminishes the activity of acetyl-CoA carboxylase in human muscle. *Diabetes* 49: 1295-1300, 2000.
5. Dzamko N, Schertzer JD, Ryall JG, Steel R, Macaulay SL, Wee S, Chen ZP, Michell BJ, Oakhill JS, Watt MJ, Jorgensen SB, Lynch GS, Kemp BE, Steinberg GR. AMPK-independent pathways regulate skeletal muscle fatty acid oxidation. *J Physiol* 586: 5819-5831, 2008.
6. Fujii N, Hayashi T, Hirshman MF, Smith JT, Habinowski SA, Kaijser L, Mu J, Ljungqvist O, Birnbaum MJ, Witters LA, Thorell A, Goodyear LJ. Exercise induces isoform-specific increase in 5'AMP-activated protein kinase activity in human skeletal muscle. *Biochem Biophys Res Commun* 273: 1150-1155, 2000.
7. Fujii N, Hirshman MF, Kane EM, Ho RC, Peter LE, Seifert MM, Goodyear LJ. AMP-activated protein kinase alpha2 activity is not essential for contraction- and hyperosmolarity-induced glucose transport in skeletal muscle. *J Biol Chem* 280: 39033-39041, 2005.
8. Hardie DG. AMP-activated/SNF1 protein kinases: conserved guardians of cellular energy. *Nat Rev Mol Cell Biol* 8: 774-785, 2007.
9. Henriquez-Olguin C, Knudsen JR, Raun SH, Li Z, Dalbram E, Treebak JT, Sylow L, Holmdahl R, Richter EA, Jaimovich E, Jensen TE. Cytosolic ROS production by NADPH oxidase 2 regulates muscle glucose uptake during exercise. *Nat Commun* 10: 4623, 2019.
10. Hoffman NJ, Parker BL, Chaudhuri R, Fisher-Wellman KH, Kleinert M, Humphrey SJ, Yang P, Holliday M, Trefely S, Fazakerley DJ, Stockli J, Burchfield JG, Jensen TE, Jothi R, Kiens B, Wojtaszewski JF, Richter EA, James DE. Global

- Phosphoproteomic Analysis of Human Skeletal Muscle Reveals a Network of Exercise-Regulated Kinases and AMPK Substrates. *Cell Metab* 22: 922-935, 2015.
11. Jensen TE, Schjerling P, Viollet B, Wojtaszewski JF, Richter EA. AMPK α 1 activation is required for stimulation of glucose uptake by twitch contraction, but not by H₂O₂, in mouse skeletal muscle. *PLoS ONE* 3: e2102, 2008.
 12. Jeppesen J, Albers PH, Rose AJ, Birk JB, Schjerling P, Dzamko N, Steinberg GR, Kiens B. Contraction-induced skeletal muscle FAT/CD36 trafficking and FA uptake is AMPK independent. *J Lipid Res* 52: 699-711, 2011.
 13. Jorgensen SB, Viollet B, Andreelli F, Frosig C, Birk JB, Schjerling P, Vaulont S, Richter EA, Wojtaszewski JF. Knockout of the α 2 but not α 1 5'-AMP-activated protein kinase isoform abolishes 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside but not contraction-induced glucose uptake in skeletal muscle. *J Biol Chem* 279: 1070-1079, 2004.
 14. Kjobsted R, Roll JLW, Jorgensen NO, Birk JB, Foretz M, Viollet B, Chadt A, Al-Hasani H, Wojtaszewski JF. AMPK and TBC1D1 Regulate Muscle Glucose Uptake After - But Not During - Exercise and Contraction. *Diabetes* 2019.
 15. Kjobsted R, Treebak JT, Fentz J, Lantier L, Viollet B, Birk JB, Schjerling P, Bjornholm M, Zierath JR, Wojtaszewski JF. Prior AICAR stimulation increases insulin sensitivity in mouse skeletal muscle in an AMPK-dependent manner. *Diabetes* 64: 2042-2055, 2015.
 16. Kurth-Kraczek EJ, Hirshman MF, Goodyear LJ, Winder WW. 5' AMP-activated protein kinase activation causes GLUT4 translocation in skeletal muscle. *Diabetes* 48: 1667-1671, 1999.
 17. Lantier L, Fentz J, Mounier R, Leclerc J, Treebak JT, Pehmoller C, Sanz N, Sakakibara I, Saint-Amand E, Rimbaud S, Maire P, Marette A, Ventura-Clapier R, Ferry A, Wojtaszewski JF, Foretz M, Viollet B. AMPK controls exercise endurance, mitochondrial oxidative capacity, and skeletal muscle integrity. *FASEB J* 28: 3211-3224, 2014.
 18. McConell GK and Kingwell BA. Does nitric oxide regulate skeletal muscle glucose uptake during exercise? *Exerc Sport Sci Rev* 34: 36-41, 2006.
 19. McConell GK, Lee-Young RS, Chen ZP, Stepto NK, Huynh NN, Stephens TJ, Canny BJ, Kemp BE. Short-term exercise training in humans reduces AMPK signalling during prolonged exercise independent of muscle glycogen. *J Physiol* 568: 665-676, 2005.
 20. McConell GK, Rattigan S, Lee-Young RS, Wadley GD, Merry TL. Skeletal muscle nitric oxide signaling and exercise: a focus on glucose metabolism. *Am J Physiol Endocrinol Metab* 303: E301-307, 2012.
 21. McGee SL, Howlett KF, Starkie RL, Cameron-Smith D, Kemp BE, Hargreaves M. Exercise increases nuclear AMPK α 2 in human skeletal muscle. *Diabetes* 52: 926-928, 2003.
 22. Merrill GF, Kurth EJ, Hardie DG, Winder WW. AICA riboside increases AMP-activated protein kinase, fatty acid oxidation, and glucose uptake in rat muscle. *Am J Physiol* 273: E1107-1112, 1997.
 23. Merry TL, Steinberg GR, Lynch GS, McConell GK. Skeletal muscle glucose uptake during contraction is regulated by nitric oxide and ROS independently of AMPK. *Am J Physiol Endocrinol Metab* 298: E577-585, 2010.

- 272 24. Mortensen B, Hingst JR, Frederiksen N, Hansen RW, Christiansen CS, Iversen N,
273 Friedrichsen M, Birk JB, Pilegaard H, Hellsten Y, Vaag A, Wojtaszewski JF. Effect
274 of birth weight and 12 weeks of exercise training on exercise-induced AMPK
275 signaling in human skeletal muscle. *Am J Physiol Endocrinol Metab* 304: E1379-
276 1390, 2013.
- 277 25. Mu J, Brozinick JT, Jr., Valladares O, Bucan M, Birnbaum MJ. A role for AMP-
278 activated protein kinase in contraction- and hypoxia- regulated glucose transport in
279 skeletal muscle. *Mol Cell* 7: 1085-1094., 2001.
- 280 26. Musi N, Fujii N, Hirshman MF, Ekberg I, Froberg S, Ljungqvist O, Thorell A,
281 Goodyear LJ. AMP-activated protein kinase (AMPK) is activated in muscle of
282 subjects with type 2 diabetes during exercise. *Diabetes* 50: 921-927, 2001.
- 283 27. Musi N, Hayashi T, Fujii N, Hirshman MF, Witters LA, Goodyear LJ. AMP-
284 activated protein kinase activity and glucose uptake in rat skeletal muscle. *Am J*
285 *Physiol Endocrinol Metab* 280: E677-684., 2001.
- 286 28. Nelson ME, Parker BL, Burchfield JG, Hoffman NJ, Needham EJ, Cooke KC,
287 Naim T, Sylow L, Ling NX, Francis D, Norris DM, Chaudhuri R, Oakhill JS,
288 Richter EA, Lynch GS, Stockli J, James DE. Phosphoproteomics reveals conserved
289 exercise-stimulated signaling and AMPK regulation of store-operated calcium
290 entry. *Embo j* e102578, 2019.
- 291 29. O'Neill HM, Lally JS, Galic S, Pulinilkunnil T, Ford RJ, Dyck JR, van Denderen
292 BJ, Kemp BE, Steinberg GR. Skeletal muscle ACC2 S212 phosphorylation is not
293 required for the control of fatty acid oxidation during exercise. *Physiol Rep* 3: 2015.
- 294 30. O'Neill HM, Lally JS, Galic S, Thomas M, Azizi PD, Fullerton MD, Smith BK,
295 Pulinilkunnil T, Chen Z, Samaan MC, Jorgensen SB, Dyck JR, Holloway GP,
296 Hawke TJ, van Denderen BJ, Kemp BE, Steinberg GR. AMPK phosphorylation of
297 ACC2 is required for skeletal muscle fatty acid oxidation and insulin sensitivity in
298 mice. *Diabetologia* 57: 1693-1702, 2014.
- 299 31. Park H, Kaushik VK, Constant S, Prentki M, Przybytkowski E, Ruderman NB,
300 Saha AK. Coordinate regulation of malonyl-CoA decarboxylase, sn-glycerol-3-
301 phosphate acyltransferase, and acetyl-CoA carboxylase by AMP-activated protein
302 kinase in rat tissues in response to exercise. *J Biol Chem* 277: 32571-32577, 2002.
- 303 32. Raney MA, Yee AJ, Todd MK, Turcotte LP. AMPK activation is not critical in the
304 regulation of muscle FA uptake and oxidation during low-intensity muscle
305 contraction. *Am J Physiol Endocrinol Metab* 288: E592-598, 2005.
- 306 33. Richter EA and Hargreaves M. Exercise, GLUT4, and skeletal muscle glucose
307 uptake. *Physiol Rev* 93: 993-1017, 2013.
- 308 34. Romijn JA, Coyle EF, Sidossis LS, Gastaldelli A, Horowitz JF, Endert E, Wolfe
309 RR. Regulation of endogenous fat and carbohydrate metabolism in relation to
310 exercise intensity and duration. *Am J Physiol* 265: E380-391, 1993.
- 311 35. Sjoberg KA, Frosig C, Kjobsted R, Sylow L, Kleinert M, Betik AC, Shaw CS,
312 Kiens B, Wojtaszewski JFP, Rattigan S, Richter EA, McConell GK. Exercise
313 Increases Human Skeletal Muscle Insulin Sensitivity via Coordinated Increases in
314 Microvascular Perfusion and Molecular Signaling. *Diabetes* 66: 1501-1510, 2017.
- 315 36. Sylow L, Jensen TE, Kleinert M, Mouatt JR, Maarbjerg SJ, Jeppesen J, Prats C,
316 Chiu TT, Boguslavsky S, Klip A, Schjerling P, Richter EA. Rac1 is a novel

- 317 regulator of contraction-stimulated glucose uptake in skeletal muscle. *Diabetes* 62:
318 1139-1151, 2013.
- 319 37. Winder WW and Hardie DG. Inactivation of acetyl-CoA carboxylase and activation
320 of AMP-activated protein kinase in muscle during exercise. *Am J Physiol* 270:
321 E299-304, 1996.
- 322 38. Wojtaszewski JF, Mourtzakis M, Hillig T, Saltin B, Pilegaard H. Dissociation of
323 AMPK activity and ACCbeta phosphorylation in human muscle during prolonged
324 exercise. *Biochem Biophys Res Commun* 298: 309-316, 2002.
- 325 39. Wojtaszewski JF, Nielsen P, Hansen BF, Richter EA, Kiens B. Isoform-specific
326 and exercise intensity-dependent activation of 5'-AMP-activated protein kinase in
327 human skeletal muscle. *J Physiol* 528: 221-226., 2000.
- 328 40. Xiao B, Sanders MJ, Underwood E, Heath R, Mayer FV, Carmena D, Jing C,
329 Walker PA, Eccleston JF, Haire LF, Saiu P, Howell SA, Aasland R, Martin SR,
330 Carling D, Gamblin SJ. Structure of mammalian AMPK and its regulation by ADP.
331 *Nature* 472: 230-233, 2011.
332